

Coronal Jets in Active Regions

Alphonse C. Sterling

Ronald L. Moore
David A. Falconer
Navdeep Panesar (NPP Postdoc)
Francisco Martinez (MSFC REU Student)
Sachiko Akiyama
Seiji Yashiro
Nat Gopalswamy

(Supported by NASA's LWS and HGI programs, NASA NPP
program, and MSFC/Hinode project.)

Overview


- ♦ Introduction
- ♦ Coronal Jets in polar coronal holes.
 - ♦ How we think they work.
- ♦ Quiet Sun jets, and how they work.
- ♦ Active region jets:
 - ♦ Example 1.
 - ♦ Example 2.
- ♦ Summary

Coronal Jets

- ♦ Well seen in X-rays (e.g., Shibata et al. 1992, Cirtain et al. 2007), and EUV (e.g., Nisticò et al. 2009).
- ♦ In polar coronal holes: size $\sim 50,000$ km x 8000 km; rate $\sim 60/\text{day}$ (Savcheva et al. 2007). Total energy $\sim 10^{26}—10^{27}$ erg (Pucci et al. 2013).
- ♦ In active regions (ARs): Similar appearance; longer ($\lesssim 10^5$ km); more energetic ($\sim 10^{27}—10^{29}$ erg; Shimojo & Shibata 2000) .
- ♦ Jets often have a “jet bright point” on one side of the jet’s base.
- ♦ Good overviews/reviews include: Shimojo et al. (1994) Nisticò et al. (2009), Raouafi et al. (2016).



Cirtain et al. (2007)





(Thanks to K. Reeves)

Minifilament Eruptions


XRT


AIA 193



Sterling et al. (Nature, 2015): 20 Polar CH jets.

“Normal” Filament Eruption (TRACE)





Sterling et al. (2015, 2016): “minifilament” eruptions.

Quiet Sun Jets - How We Think They Work:

Answer: The same as polar coronal hole jets!

(Panesar et al. 2016, ApJL; 10 quiet Sun jets)

AIA 171


AIA 94




Active Region Jets: Example 1

- Single AR, AR 11513
- Only ~8-hr period, but many jets.
- AIA+HMI; no Hinode, but some SXI images.
- Sterling et al. (2016), ApJ


AIA 171



Sterling et al. (2016, ApJ)




AIA 304




AIA 94



AIA 304




AIA 94




AR Jet Example 2: To investigate further, look at a different AR:


- 14 Jan 2015 (NOAA AR 12259).
- AIA, HMI, Hinode, IRIS





Hinode/XRT


AIA 193







HMI of IRIS-observed region:



Jets occur at *flux cancelation locations!

For this jet:

- This jet (actually, a series of jets) is consistent with the minifilament-eruption schematic.
- Minifilament itself however is not as clear as in other cases.
- Occurs at neutral line.
- Cancellation at the neutral line.
(Strength? Gradient?).

Flux Cancellation Rates:

Preliminary values (Panesar et al. 2016; Sterling et al., in preparation):

- For quiet Sun jets: $(1.7 \pm 0.7) \times 10^{18}$ erg
- For AR jets: $(0.5 \pm 1.0) \times 10^{19}$ erg

Summary and Conclusions

- ♦ Detailed investigations of several AR jets.
- ♦ Among those we investigated:
 - ♦ Visually all fit the minifilament picture, from the magnetic-field setup standpoint.
 - ♦ Slower-buildup ones have obvious minifilaments.
 - ♦ Faster-buildup ones may have minifilaments, but of a different character than what we're used to (thinner, and maybe hidden by emission). Or maybe a different process.
- ♦ All occur on neutral lines.
- ♦ Frequently (if not always) there is flux cancelation occurring on the neutral lines. There are definitely cases of emergence+cancelation, and there may be cases of minifilament eruption triggered by emerging flux. (Cf. Mulay et al. 2016, others....)